



Individual Susceptibility to Hypobaric Environments: An Update

Jennifer Law, M.D.
University of California Davis Medical Center
Department of Emergency Medicine

Sharmi Watkins, M.D., M.P.H.
Element Scientist, Exploration Medical Capability
The University of Texas Medical Branch
NASA Johnson Space Center Bioastronautics Contract

THE NASA STI PROGRAM OFFICE . . . IN PROFILE

Since its founding, NASA has been dedicated to the advancement of aeronautics and space science. The NASA Scientific and Technical Information (STI) Program Office plays a key part in helping NASA maintain this important role.

The NASA STI Program Office is operated by Langley Research Center, the lead center for NASA's scientific and technical information. The NASA STI Program Office provides access to the NASA STI Database, the largest collection of aeronautical and space science STI in the world. The Program Office is also NASA's institutional mechanism for disseminating the results of its research and development activities. These results are published by NASA in the NASA STI Report Series, which includes the following report types:

- **TECHNICAL PUBLICATION.** Reports of completed research or a major significant phase of research that present the results of NASA programs and include extensive data or theoretical analysis. Includes compilations of significant scientific and technical data and information deemed to be of continuing reference value. NASA's counterpart of peer-reviewed formal professional papers but has less stringent limitations on manuscript length and extent of graphic presentations.
- **TECHNICAL MEMORANDUM.** Scientific and technical findings that are preliminary or of specialized interest, e.g., quick release reports, working papers, and bibliographies that contain minimal annotation. Does not contain extensive analysis.
- **CONTRACTOR REPORT.** Scientific and technical findings by NASA-sponsored contractors and grantees.
- **CONFERENCE PUBLICATION.** Collected papers from scientific and technical conferences,

symposia, seminars, or other meetings sponsored or cosponsored by NASA.

- **SPECIAL PUBLICATION.** Scientific, technical, or historical information from NASA programs, projects, and mission, often concerned with subjects having substantial public interest.
- **TECHNICAL TRANSLATION.** English-language translations of foreign scientific and technical material pertinent to NASA's mission.

Specialized services that complement the STI Program Office's diverse offerings include creating custom thesauri, building customized databases, organizing and publishing research results . . . even providing videos.

For more information about the NASA STI Program Office, see the following:

- Access the NASA STI Program Home Page at <http://www.sti.nasa.gov>
- E-mail your question via the internet to help@sti.nasa.gov
- Fax your question to the NASA Access Help Desk at (301) 621-0134
- Telephone the NASA Access Help Desk at (301) 621-0390
- Write to:

NASA Access Help Desk

NASA Center for AeroSpace Information

7115 Standard

Hanover, MD 21076-1320



Individual Susceptibility to Hypobaric Environments: An Update

Jennifer Law, M.D.
University of California Davis Medical Center
Department of Emergency Medicine

Sharmi Watkins, M.D., M.P.H.
Element Scientist, Exploration Medical Capability
The University of Texas Medical Branch
NASA Johnson Space Center Bioastronautics Contract

Available from:

NASA Center for Aerospace Information
7115 Standard Drive
Hanover, MD 21076-1320
301-621-0390

National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161
703-605-6000

This report is also available in electronic form at <http://ston.jsc.nasa.gov/collections/TRS/>

ABSTRACT

Astronauts are at risk for developing decompression sickness (DCS) while exposed to the hypobaric environment of the extravehicular suit in space, in terrestrial hypobaric chambers, and during ascent from neutral buoyancy training dives. There is increasing recognition that DCS risk is different between diving and altitude exposures, with many individual parameters and environmental factors implicated as risk factors for development of DCS in divers but are not recognized as risk factors in altitude exposures. Much of the literature to date has focused on patent foramen ovale (PFO), which has long been considered a major risk factor for DCS in diving exposures, but its link to serious DCS in altitude exposures remains unclear. Knowledge of those risk factors specific to hypobaric DCS may help identify susceptible individuals and aid in astronaut selection, crew assignment, and mission planning. This paper reviews the current literature pertaining to these risk factors, including PFO, anthropometric parameters, gender, menstrual cycle, lifetime diving experience, physical fitness, biochemical levels, complement activation, cigarette smoking, fluid balance, and ambient temperature. Further research to evaluate pertinent risk factors for DCS in altitude exposures is recommended.

Contents

Introduction	1
Patent Foramen Ovale & Decompression Sickness	1
PFO Committee/Medical Operations EVA IPT	2
Beyond the Med Ops EVA IPT Report	3
Other Factors Affecting Individual Susceptibility to DCS	4
Individual Parameters	4
Environmental Factors	6
Conclusion.....	6
References	7

Introduction

Absolute or relative reductions in ambient pressure, as experienced during extravehicular activity, physiological training in hypobaric chambers, or ascent from diving, can all place unique stressors on the human body. Participants in these environments are at risk for the development of decompression sickness (DCS), whose symptoms range from mild musculoskeletal pain to serious neurological deficits and even to death. Given the potential negative impact on crew health and mission success, it would be invaluable to identify risk factors for DCS and employ preventive strategies to mitigate the risk.

DCS occurs as a result of transitioning from one environment to a more hypobaric environment. When the ambient pressure decreases, nitrogen dissolved in the bloodstream comes out of solution and forms bubbles, which may remain locally or may be transported into the venous system. Most bubbles are trapped in the pulmonary capillary bed before being quickly resorbed, but if the pulmonary filter gets overwhelmed or bypassed due to an anatomic defect allowing for right-to-left shunt, these bubbles may become arterial gas emboli and result in local ischemia anywhere in the body.

While Type I DCS (“pain only” or mild lymphatic involvement) generally requires no treatment or brief treatment with 100% oxygen, Type II DCS can manifest with pulmonary, cardiovascular, or neurologic symptoms that are more difficult to treat and may result in significant morbidity or mortality.¹ The Human Research Program’s Exploration Medical Capability Element has been tasked with reviewing what is known about individual susceptibility to hypobaric environments—specifically, DCS. This white paper will summarize the work done by two sub-groups in 1999, the Patent Foramen Ovale Committee and Medical Operations EVA Integrated Product Team. The paper will also review the literature on PFO since 1999, and discuss other risk factors for DCS that have been proposed.

Patent Foramen Ovale & Decompression Sickness

Patent foramen ovale (PFO), the primary means of arterial gas embolism through the heart, has been implicated in an individual’s susceptibility to DCS. PFO is a common remnant of fetal circulation when the atrial septum primum and septum secundum fail to fuse during the first months of life, enabling abnormal blood flow between the left and right atria. Some PFOs open only when there is an increase in pulmonary pressure—e.g., during a Valsalva maneuver or strenuous activity—but some PFOs are persistently open and believed to be more dangerous since they pose a constant threat for arterializations of gas bubbles.² An autopsy study reported that the overall incidence of PFO in 965 subjects was 27.3% with a median diameter of 5 mm; 9.7% of the subjects had a PFO greater than 5 mm.³ The incidence of resting PFO was reported to be 5 to 10% in the literature and might be as high as 21% in the experience of the JSC Cardiovascular Laboratory.⁴

Concern about the role of PFO in the development of decompression sickness arose from diving data. Studies have shown that divers with a PFO have a 2.5 to 4.5 fold increase in the risk of developing type II DCS.⁵⁻⁶ To date, PFO has been the most studied risk factor of DCS, although its role in altitude exposures remains controversial for a number of reasons, including the lack of a standardized technique for detecting PFOs and uncertainty about the exact incidence of DCS.

PFO Committee/Medical Operations EVA IPT

In 1999, a NASA-sponsored committee was tasked with reviewing data on PFO and making recommendations about the merit of PFO screening in the astronauts and test subjects. The Medical Operations EVA Integrated Product Team (Med Ops EVA IPT) subsequently reviewed the committee's recommendations and consulted outside experts to develop a consensus position on PFO screening.

Despite the strength of evidence in diving exposures, PFO had not been directly linked to Type II DCS in altitude exposures. A retrospective study by the Navy found no association between PFO and Type II DCS in 45 aviators who presented with Type II DCS symptoms related to altitude physiological training.⁷

In an Air Force altitude chamber-based study in which 1,500 volunteer subjects were monitored for gas emboli simultaneously in the right and left sides of the heart at simulated altitude, six subjects were found to have right-to-left gas crossover; five of these subjects were symptomatic with joint pain or skin mottling (but no cerebral symptoms) at the time of arterial gas embolism. Of the six cases, one was found to have a PFO, one had a small sinus venosus defect, three had no septal defect, and one was not available for evaluation.⁸ In another presentation, Pilmanis reported 39 cases of Type II DCS in over 2,000 research exposures which simulated EVA exposures of 4-hour durations or longer. None of these subjects had detectable gas phase crossover during Trans Thoracic Echocardiography (TTE) monitoring of all four cardiac chambers.

While no published study showed a statistically significant association between PFO and Type II DCS in altitude exposures, case reports did suggest a relationship. Past hypobaric chamber research at JSC identified five cases of Type II DCS. Three of these cases were tested: two (66%) were found to have a PFO. Of note, four of the five cases did not prebreathe oxygen. A serious case of Type II DCS was reported at Duke University during testing of a prebreathe reduction program, despite two hours of oxygen prebreathe and albeit a less conservative nitrogen elimination protocol than the ISS operational protocol. This subject was later determined to have a resting PFO.

Based on the available data, the PFO Committee recommended screening all individuals for high-risk PFO, such as resting PFO, PFO in excess of 5 mm in diameter, or individuals with more than 20 countable bubbles observed passing through the PFO during a measurement. It was noted, however, that some PFOs that opened only with provocation during strenuous EVA, might have a greater diameter than a smaller resting PFO. Furthermore, TTE with bubble contrast, the screening technique recommended by the PFO Committee, had a sensitivity of 60 to 70%, and identifying "high risk PFOs" was technically challenging and somewhat subjective. Transesophageal echocardiography was considered the gold standard but invasive and not without risk.

The Med Ops EVA IPT noted that PFO screening could potentially eliminate 10 to 25% of EVA-proficient crewmembers from the existing cadre of astronauts, which would severely limit the operational flexibility of assigning crewmembers to station missions. Furthermore, given the limitations of PFO screening as well as the unclear relationship between PFO and Type II DCS in altitude exposures, PFO screening would not eliminate the risk of Type II DCS on orbit. A more effective means of mitigating the risk of Type II DCS would be to minimize the chance of bubble formation by oxygen prebreathe procedures.

Taking the PFO Committee's recommendations and external experts' review into account, the Med Ops EVA IPT recommended the following:

1. That NASA continue to eliminate astronaut candidates with flow-significant atrial septal defects detected with TTE
2. That an in-suit Doppler be implemented as soon as possible, and that an on-orbit operational validation of the 2-hour prebreathe protocol be conducted
3. That a multi-center retrospective study be initiated to determine the association between PFO and Type II DCS in well-controlled altitude exposures, as a function of decompression stress
4. That other prospective PFO studies be considered for funding
5. That the possibility of screening out astronauts for high risk PFO continue to be considered as warranted by new data or diagnostic modalities
6. That sensitive screening methods for the risk of Type II DCS be developed, beyond screening for PFO, i.e., screening for a combination of PFO and the propensity to produce high-grade venous gas embolism, which would occur in a very small percentage of astronauts

Beyond the Med Ops EVA IPT Report

Since 1999, other retrospective studies and meta-analyses have supported the relationship between PFO and Type II DCS in divers. In a retrospective cohort study of 52 sport divers, Schwerzmann et al.⁹ found a 4.5-fold increased risk for DCS in divers with PFO than those without a PFO. On MRI, almost twice as many ischemic brain lesions were seen in divers with PFO, but the difference was not statistically significant. The authors did not analyze the relationship between these brain lesions and DCS. Gempp et al.¹⁰ studied 49 divers with spinal cord DCS and found that divers with DCS were 3.6 times more likely than healthy divers to have a large right-to-left shunt on transcranial Doppler ultrasonography. While shunting was not associated with an increased incidence of cervical spinal cord DCS, divers with a right-to-left shunt were 6.9 times more likely to have thoracolumbar DCS. These studies confirm the increased prevalence of PFO or right-to-left shunting in divers who experience DCS and begin to offer a neuropathological basis for symptoms seen in Type II DCS.

New data also suggest that PFOs evolve as divers age. Anecdotally, some divers suddenly become susceptible to DCS after an uneventful diving career of many years. A longitudinal study spanning 6 to 8 years found that the permeability of the foramen ovale in 40 divers changed. A small number (7.5%) of

the divers who initially had a PFO actually saw closure of the PFO during this study. Another 22.5% of the study cohort developed increased permeability, half of whom had no initial PFO.¹¹

In altitude exposures, it remains unclear whether PFO is a risk factor for Type II DCS. An evidence-based literature review concluded that PFO could occur in altitude-induced DCS, but there was no information beyond two case series to indicate causality.¹² Under-reporting of symptoms in altitude conditions due to perceived negative career impact might partly explain the paucity of data.

It has been suggested that diving and altitude exposures are fundamentally different entities, as evidenced by the observation that the most common symptoms of DCS at altitude are pain only, as opposed to neurologic symptoms seen in recreational divers¹³; moreover, DCS symptoms tend to occur during altitude exposure versus after diving exposure.¹⁴ Differences in duration of exposure, prebreathe protocols, and oxygen use during exposure, as in the case of EVA, may further account for the difference in diving and altitude data. Review of the literature since 1999 reveals that additional research is needed to ascertain the relationship between PFO and Type II DCS in altitude exposures. Thus, the benefit of screening astronauts for PFO remains unclear.

Other Factors Affecting Individual Susceptibility to DCS

While patent foramen ovale has been studied the most, other risk factors for DCS have been proposed based on known mechanisms, including bubble crossover through the heart, bubble crossover through the lungs, and autochthonous bubbles. PFO and right-to-left shunting represent only the first mechanism. Little is known about conditions that allow crossover through the lungs, which appear to have a threshold for filtration of venous bubbles by the pulmonary vasculature, at least in dogs.¹⁵ A variety of individual parameters and environmental factors have been studied in an effort to identify risk factors for autochthonous bubble formation that lead to susceptibility to DCS.

Individual Parameters

Parameters such as age, height, weight, body mass index, percent body fat, aerobic capacity, and gender have been reported to influence DCS incidence.¹⁶⁻¹⁹ In the most comprehensive study of these parameters to date, Air Force researchers sought to quantify individual susceptibility to DCS and determine if the aforementioned variables could be used to predict DCS risk. Retrospectively reviewing 2,980 altitude exposures in the Air Force Laboratory Altitude DCS Research Database,²⁰ a study found that the combination of lower VO_{2max} and greater weight appeared to be the best predictor of DCS, but this accounted for less than 13% of the variation in DCS susceptibility. The authors concluded that individual susceptibility to DCS could not be predicted by these anthropometric and physiologic variables.

In another study, Webb et al.¹⁷ prospectively compared male and female subjects in 961 exposures to simulated altitude and found no statistically significant difference in DCS incidence between men (49.5%) and women (45.3%). Among women, they found no difference in DCS during the first two weeks of the menstrual cycle between users of hormonal contraception vs. non-users. However, if only

the last half of the menstrual cycle were considered, hormonal contraception appeared to double the risk of DCS, a finding supporting an earlier report by Doyle et al.²¹ but refuted by Lee et al.²²

Lee and her colleagues surveyed 240 female sports divers treated with hyperbaric therapy for DCS in 23 treatment centers worldwide, and found the incidence of all DCS symptoms to be greatest in the first half of the menstrual cycle (29 to 34%), with a marked fall in the third week (13 to 23%) before rising again in the fourth week (21 to 25%); the trend was similar for Type I and Type II symptoms. The effect of contraceptive use appeared to shift the occurrence of DCS to later in the cycle but there was no significant difference between contraceptive users and non-users. There was, however, a statistically significant difference between the two groups when age was taken into account. The observed differences in these studies might have been due to differences in how menstrual history was recorded as well as doses and formulation of contraceptives. More study is needed to draw a conclusion about the role of menstruation and contraceptive use in the risk of DCS in women, and whether EVA schedules would need to be adjusted accordingly.

An individual's lifetime diving experience has also been linked to DCS risk. In a survey-based study of 429 subjects, Klingmann et al.²³ found a 2.51-fold higher lifetime incidence of DCI in divers with a history of deep dives greater than 40 meters compared to divers whose lifetime maximal depth was 40 meters or less. The relative risk of DCS between divers with 200 or fewer dives and "highly experienced" divers with more than 400 dives was 3.90. Similarly, certification to lower depths was associated with 1.97 to 8.17 times higher incidence of DCS. There was no statistically significant difference between technical divers and non-technical divers. Attrition, rather than any physiologic factors, was identified as the explanation for the association between low-depth diving experience and risk of DCS, but the authors did acknowledge a selection bias in their methodology and called for further study before their results could be generalized to all divers. Similarly, prior DCS might predispose an individual to subsequent development of DCS, but data have been limited to observations of caisson workers in the 1970s.²⁴

On the other hand, a more recent study suggested that repeated scuba dives and regular physical exercise activity might protect against DCS, possibly due to reduced bubble formation.²⁵ However, exercise itself has been observed to offer protection from DCS. In addition to laboratory studies that showed sedentary rats and pigs were more susceptible to severe DCS and death than exercise conditioned groups,²⁶⁻²⁷ several human studies have found decreased venous gas emboli formation, thereby inferring decreased DCS risk, in divers who exercised 24 hours before diving,²⁸ 2 hours before diving,²⁹ during decompression,³⁰ and during a decompression stop.³¹ In altitude exposures, exercise during prebreathe also appeared to be beneficial in preventing DCS.^{19,32}

Biochemically, limited studies have suggested that subjects susceptible to the formation of venous gas emboli during decompression had significant higher levels of total cholesterol, high-density lipoprotein cholesterol, potassium, phosphate, calcium, and magnesium.³³ A follow-on study was unable to replicate these findings, and it was noted that factors such as prebreathe time, use of an intermediate pressure stage, and length of time in the chamber could affect various biochemical parameters.³⁴

The observation that many of the effects of complement activation were similar to the symptoms of DCS led to several studies that ultimately disproved complement activation as a risk factor for DCS. Ward et al.³⁵ found that the presence of air bubbles in plasma activated the complement system by the alternate pathway and that subjects who were more sensitive to this complement activation were more susceptible to DCS. However, in vivo testing failed to demonstrate this phenomenon, and subjects who developed DCS showed no difference in complement activation during dives compared to healthy subjects.³⁶

Finally, a history of cigarette smoking may predispose an individual to DCS. A retrospective analysis of the Divers Alert Network (DAN) database found that heavy smokers (>15 pack-year history) were 1.88 times more likely to have DCS than divers who had never smoked and 1.56 times more likely than light smokers (0-15 pack-year history).³⁷

Environmental Factors

Fluid balance appears to play a role in DCS. One study randomized swine subjects into two groups: a hydrated group that was allowed ad lib access to water during a simulated saturation dive, and a dehydrated group that was given intravenous furosemide without access to water. The dehydrated group showed a significantly increased rate and faster onset of DCS and death, possibly due to increased bubble formation or altered nitrogen removal.³⁸

Ambient temperature has also been considered as a risk factor for DCS. A Navy study sought to compare the incidence of DCS in divers who were immersed in air decompression dives to 120 feet of seawater at either 97°F or 80°F, and found that warm conditions during bottom time and cold conditions during decompression increased the risk of DCS, but this finding was confounded by dehydration.³⁹ Other studies found that hot water suits were 1.81 to 1.96 times more likely than passive thermal protection to be associated with DCS, although some of these findings might have been attributable to the dive profile or bottom time.⁴⁰ It has been proposed that increased susceptibility to DCS may result from increased gas uptake in warm conditions and decreased elimination in cold conditions, but there is insufficient data to confirm this theory.⁴¹

Conclusion

The ability to predict an individual's susceptibility to decompression sickness in hypobaric environments would be advantageous for astronaut selection, crew assignment, and mission planning. To date, much focus has been on PFO as a risk factor for DCS, but a number of other individual parameters as well as environmental factors have also been implicated in increased risk of DCS. Although diving exposures were once considered analogous to altitude exposures, there is increasing appreciation for the fundamental differences between diving and altitude. There continues to be a need for altitude research under controlled conditions to evaluate potential risk factors for DCS in altitude exposures and apply knowledge about individual susceptibilities to mission planning. In the meantime, it seems sensible to continue current protective measures such as exercise-enhanced prebreathe periods to minimize the risk of DCS in hypobaric environments.

References

- 1 Francis TJR, Dutka AJ, Hallenbeck JM. Pathophysiology of decompression sickness. In: Bove AA, Davis JC, ed. *Diving Medicine*, Philadelphia: WB Saunders, 1990:170-87.
- 2 Medical Operations EVA IPT. Recommendations Regarding PFO Screening. Internal Report, 1999.
- 3 Hagen PT, Scholz DG, Edwards WD. Incidence and size of patent foramen ovale during the first 10 decades of life: an autopsy study of 965 normal hearts. *Mayo Clin Proc.* 1984 Jan;59(1):17-20.
- 4 Waligora JM, Powell MR, Norfleet WT, Wood M, Martin D. The incidence of patent foramen ovale and serious symptoms of decompression sickness in trials conducted to develop operational decompression protocols for space flight. Internal Report, 1999.
- 5 Bove AA. Risk of decompression sickness with patent foramen ovale. *Undersea Hyper Med.* 1998;25:175-78.
- 6 Drighil A, El Mosalami H, Elbadaoui N, Chraibi S, Bennis A. Patent foramen ovale: a new disease? *Int J Cardiol.* 2007 Oct 31;122(1):1-9.
- 7 Gallagher KL, Hopkins EW, Clark JB, Hawley TA. US Navy experience with Type II decompression sickness and the association with patent foramen ovale. *Aviat Space Environ Med.*, abstract #290, 1996;67:712.
- 8 Pilmanis AA, Meissner FW, Olson RM. Left ventricular gas emboli in six cases of altitude-induced decompression sickness. *Aviat Space Environ Med.* 1996 Nov;67(11):1092-6.
- 9 Schwerzmann M, Seiler C, Lipp E, Guzman R, Lövblad KO, Kraus M, Kucher N. Relation between directly detected patent foramen ovale and ischemic brain lesions in sport divers. *Ann Intern Med.* 2001 Jan 2;134(1):21-4.
- 10 Gempp E, Blatteau JE, Stephant E, Louge P. Relation between right-to-left shunts and spinal cord decompression sickness in divers. *Int J Sports Med.* 2009 Feb;30(2):150-3.
- 11 Germonpré P. Patent foramen ovale and diving. *Cardiol Clin.* 2005 Feb;23(1):97-104.
- 12 Straus SE, Sackett DL. Evidence-based literature review of the relation between patent foramen ovale (PFO) and decompression sickness (DCS). Report to the Canadian Space Agency, 2001.
- 13 Moon RE. Patent foramen ovale (PFO) and decompression illness (DCI) in space. *Aviat Space Environ Med.*, abstract #380, 2000; 71(3):352-8.

- 14 Pilmanis AA, Petropoulos LJ, Kannan N, Webb JT. Decompression sickness risk model: development and validation by 150 prospective hypobaric exposures. *Aviat Space Environ Med.* 2004 Sep;75(9):749-59.
- 15 Butler BD, Hills BA. Transpulmonary passage of venous air emboli. *J Appl Physiol.* 1985 Aug;59(2):543-7.
- 16 Sulaiman ZM, Pilmanis AA, O'Connor RB. Relationship between age and susceptibility to altitude decompression sickness. *Aviat Space Environ Med.* 1997 Aug;68(8):695-8.
- 17 Conkin J, Powell MR, Gernhardt ML. Age affects severity of venous gas emboli on decompression from 14.7 to 4.3 psia. *Aviat Space Environ Med.* 2003 Nov;74(11):1142-50.
- 18 Webb JT, Kannan N, Pilmanis AA. Gender not a factor for altitude decompression sickness risk. *Aviat Space Environ Med.* 2003 Jan;74(1):2-10.
- 19 Webb JT, Pilmanis AA, Buldin UI. Altitude decompression sickness at 7620 m following prebreathe enhanced with exercise periods. *Aviat Space Environ Med.* 2004; 75(10):859-64.
- 20 Webb JT, Pilmanis AA, Balldin UI, Fischer JR. Altitude decompression sickness susceptibility: influence of anthropometric and physiologic variables. *Aviat Space Environ Med.* 2005 Jun;76(6):547-51.
- 21 Doyle K, Baek PS, De Long ER, et al. Menstruation as a risk factor for decompression illness (DCI) in female scuba divers taking oral contraceptives (OC). *Undersea Hyperb Med* 1997; 24(Suppl):A143.
- 22 Lee V, St Leger Dowse M, Edge C, Gunby A, Bryson P. Decompression sickness in women: a possible relationship with the menstrual cycle. *Aviat Space Environ Med.* 2003 Nov;74(11):1177-82.
- 23 Klingmann C, Gonnermann A, Dreyhaupt J, Vent J, Praetorius M, Plinkert PK. Decompression illness reported in a survey of 429 recreational divers. *Aviat Space Environ Med.* 2008 Feb;79(2):123-8.
- 24 Francis TJR, Mitchell RA. Pathophysiology of decompression sickness. In: Brubakk A, Neuman T, ed. *Bennett and Elliott's Physiology and Medicine of Diving*, Philadelphia: WB Saunders, 2002:544-546.
- 25 Pontier JM, Guerrero F, Castagna O. Bubble formation and endothelial function before and after 3 months of dive training. *Aviat Space Environ Med.* 2009 Jan;80(1):15-9.
- 26 Broome JR, Dutka AJ, McNamee GA. Exercise conditioning reduces the risk of neurologic decompression illness in swine. *Undersea Hyperb Med.* 1995 Mar;22(1):73-85.

- 27 Wisløff U, Brubakk AO. Aerobic endurance training reduces bubble formation and increases survival in rats exposed to hyperbaric pressure. *J Physiol*. 2001 Dec 1;537(Pt 2):607-11.
- 28 Đujić Z, Duplancic D, Marinovic-Terzic I, Bakovic D, Ivancev V, Valic Z, Eterovic D, Petri NM, Wisløff U, Brubakk AO. Aerobic exercise before diving reduces venous gas bubble formation in humans. *J Physiol*. 2004 Mar 16;555(Pt 3):637-42.
- 29 Blatteau JE, Gempp E, Galland FM, Pontier JM, Sainty JM, Robinet C. Aerobic exercise 2 hours before a dive to 30 msw decreases bubble formation after decompression. *Aviat Space Environ Med*. 2005;76:666-9.
- 30 Jankowski LW, Nishi RY, Eaton DJ, Griffin AP. Exercise during decompression reduces the amount of venous gas emboli. *Undersea Hyperb Med*. 1997 Jun;24(2):59-65.
- 31 Đujić Z, Palada I, Obad A, Duplancić D, Baković D, Valic Z. Exercise during a 3-min decompression stop reduces postdive venous gas bubbles. *Med Sci Sports Exerc*. 2005 Aug;37(8):1319-23.
- 32 Gernhardt ML, Conkin J, Foster PP, et al. Design and testing of a two hour oxygen prebreathe protocol for space walks from the International Space Station. *Undersea Biomed Res* 2000; 27:12.
- 33 Jauchem JR, Waligora JM, Conkin J, Horrigan DJ Jr, Johnson PC Jr. Blood biochemical factors in humans resistant and susceptible to formation of venous gas emboli during decompression. *Eur J Appl Physiol Occup Physiol*. 1986;55(1):68-73.
- 34 Jauchem JR, Waligora JM, Johnson PC Jr. Blood biochemical and cellular changes during decompression and simulated extravehicular activity. *Int Arch Occup Environ Health*. 1990;62(5):391-6.
- 35 Ward CA, McCullough D, Fraser WD. Relation between complement activation and susceptibility to decompression sickness. *J Appl Physiol*. 1987 Mar;62(3):1160-6.
- 36 Shastri KA, Logue GL, Lundgren CE, Logue CJ, Suggs DF. Diving decompression fails to activate complement. *Undersea Hyperb Med*. 1997 Jun;24(2):51-7.
- 37 Buch DA, El Moalem H, Dovenbarger JA, Uguccioni DM, Moon RE. Cigarette smoking and decompression illness severity: a retrospective study in recreational divers. *Aviat Space Environ Med*. 2003;74:1271-4.
- 38 Fahlman A, Dromsky DM. Dehydration effects on the risk of severe decompression sickness in a swine model. *Aviat Space Environ Med* 2006; 77:102-6.
- 39 Gerth WA, Ruterbusch VL, Long ET. The influence of thermal exposure on diver susceptibility to decompression sickness. Navy Report NEDU TR 06-07, 2007.

- 40 Leffler, CT. Effect of ambient temperature on the risk of decompression sickness in surface decompression divers. *Aviat Space Environ Med.* 2001 May;72(5):477-483.
- 41 Toner CB, Ball R. The effect of temperature on decompression and decompression sickness risk: a critical review. *Naval Medical Research Institute Report NMRC 2004-003*, 2004.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave Blank)	2. REPORT DATE May 2010	3. REPORT TYPE AND DATES COVERED NASA Technical Paper		
4. TITLE AND SUBTITLE Individual Susceptibility to Hypobaric Environments: An Update			5. FUNDING NUMBERS	
6. AUTHOR(S) Jennier Law, Sharmi Watkins				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Lyndon B. Johnson Space Center Houston, Texas 77058			8. PERFORMING ORGANIZATION REPORT NUMBERS S-1068	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Washington, DC 20546-0001			10. SPONSORING/MONITORING AGENCY REPORT NUMBER TP-2010-216123	
11. SUPPLEMENTARY NOTES NASA Johnson Space Center				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Available from the NASA Center for AeroSpace Information (CASI) 7115 Standard Hanover, MD 21076-1320 Category: 52			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Astronauts are at risk for developing decompression sickness (DCS) while exposed to the hypobaric environment of the extravehicular suit in space, in terrestrial hypobaric chambers, and during ascent from neutral buoyancy training dives. There is increasing recognition that DCS risk is different between diving and altitude exposures, with many individual parameters and environmental factors implicated as risk factors for development of DCS in divers but are not recognized as risk factors in altitude exposures. Much of the literature to date has focused on patent foramen ovale (PFO), which has long been considered a major risk factor for DCS in diving exposures, but its link to serious DCS in altitude exposures remains unclear. Knowledge of those risk factors specific to hypobaric DCS may help identify susceptible individuals and aid in astronaut selection, crew assignment, and mission planning. This paper reviews the current literature pertaining to these risk factors, including PFO, anthropometric parameters, gender, menstrual cycle, lifetime diving experience, physical fitness, biochemical levels, complement activation, cigarette smoking, fluid balance, and ambient temperature. Further research to evaluate pertinent risk factors for DCS in altitude exposures is recommended.				
14. SUBJECT TERMS decompression sickness, DCS, hypobaric, patent foramen ovale, PFO			15. NUMBER OF PAGES 20	16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	
